



www.ofthalmoloji.org

TURKISH JOURNAL OF OPHTHALMOLOGY

TURKISH JOURNAL OF OPHTHALMOLOGY

TJO

E-ISSN: 2149-8709

Research Articles

Comparison of 20% Autologous Platelet-Rich Plasma Versus Conventional Treatment in Moderate to Severe Dry Eye Patients

Sachan et al.; Prayagraj, India

Comparative Analysis of Automated vs. Expert-Designed Machine Learning Models in Age-Related Macular Degeneration Detection and Classification

Durmaz Engin et al.; İzmir, Türkiye

Pattern of RNFL Damage in Early- and Late-Stage Primary Open-Angle Glaucoma Using the Disc Damage Likelihood Scale and Optical Coherence Tomography

Ülgin et al.; Lahore, Pakistan

Real-Life Effectiveness and Safety of Selective Laser Trabeculoplasty as Primary, Adjunctive, and Substitutive Therapy

Oliver-Gutierrez et al.; Barcelona, Spain

Gonioscopy-Assisted Transluminal Trabeculotomy versus Bent Ab Interno Needle Goniectomy in Patients with Open-Angle Glaucoma

Üçgül et al.; Kırşehir, Ankara, Türkiye

Reviews

Binocular Approaches in Amblyopia Treatment Based on Dichoptic Stimulation

Yabanoğlu and Taylan Şekeroğlu; Ankara, Türkiye

Ab Interno Goniectomy/Goniectomy Techniques

Aktaş et al.; San Antonio, TX, Jacksonville, FL, St Louis, MO, Lexington, KY, Miami, FL, USA; Abu Dhabi, UAE; Kırşehir, Türkiye

Letters to the Editor

A Promising Outcome of the Augmented Modified Hummelsheim Procedure in a Challenging Case of Inferior Rectus Hypoplasia

Priscilia and Bani; Jakarta, Indonesia

Ptosis Repair by "PEANUTS" MMCR: "Pelin's Easy and Needle Up To Stretch" Müller's Muscle Conjunctival Resection Without the Putterman Clamp

Kaynak et al.; İstanbul, İzmir, Türkiye

May
June 2025
55 Volume
Issue 3

TURKISH JOURNAL OF OPHTHALMOLOGY



www.offtalmoloji.org

TJO

Editor-in-Chief

Banu BOZKURT, MD

Selçuk University Faculty of Medicine, Department of Ophthalmology, Konya, Türkiye

Areas of Interest: Cornea and Ocular Surface Disease, Glaucoma, Allergy and Immunology, Contact Lens

E-mail: drbanubozkurt@yahoo.com

ORCID ID: orcid.org/0000-0002-9847-3521

Associate Editors

Sait EĞRİLMEZ, MD

İzmir University of Economics Faculty of Medicine, İzmir, Türkiye

Areas of Interest: Cornea and Ocular Surface Disease, Contact Lens, Refraction, Cataract and Refractive Surgery

E-mail: saitegrilmez@gmail.com

ORCID ID: orcid.org/0000-0002-6971-527X

Hakan ÖZDEMİR, MD

Bezmialem Vakıf University Faculty of Medicine, Department of Ophthalmology, İstanbul, Türkiye

Areas of Interest: Medical Retina, Vitreoretinal Surgery

E-mail: hozdemir72@hotmail.com

ORCID ID: orcid.org/0000-0002-1719-4265

Nilgün YILDIRIM, MD

Eskişehir Osmangazi University Faculty of Medicine, Department of Ophthalmology, Eskişehir, Türkiye

Areas of Interest: Glaucoma, Cornea and Ocular Surface, Oculoplastic Surgery

E-mail: nyildirim@yahoo.com

ORCID ID: orcid.org/0000-0001-6506-0336

Özlem YILDIRIM, MD

Mersin University Faculty of Medicine, Department of Ophthalmology, Mersin, Türkiye

Areas of Interest: Uveitis, Medical Retina, Glaucoma

E-mail: dryildirimoz@hotmail.com

ORCID ID: orcid.org/0000-0002-3773-2497

Statistics Editor

Ahmet DİRİCAN,

İstanbul University İstanbul Faculty of Medicine, Department of Biostatistics and Medical Informatics, İstanbul, Türkiye

English Language Editor

Jacqueline Renee GUTENKUNST, MARYLAND, USA

Advisory Board

Özgül ALTINTAŞ,

Acıbadem Maslak Hospital, Clinic of Ophthalmology, Private Practice, İstanbul, Türkiye

Halil Özgür ARTUNAY,

University of Health Sciences Türkiye, Beyoğlu Eye Training and Research Hospital, Clinic of Ophthalmology, İstanbul, Türkiye

Jose M. BENÍTEZ-del-CASTILLO,

Universidad Complutense de Madrid, Hospital Clínico San Carlos, Department of Ophthalmology, Madrid, Spain

Ayşe BURCU,

University of Health Sciences Türkiye, Ankara Training and Research Hospital, Clinic of Ophthalmology, Ankara, Türkiye

Virginia CALDER,

UCL Institute of Ophthalmology, Department of Ocular Immunology, London, UK

Doğan CEYHAN,

Güven Hospital Çayyolu Medical Center, Clinic of Ophthalmology, Ankara, Türkiye

M. Pınar ÇAKAR ÖZDAL,

Private Practice, Ankara, Türkiye

Ebru Nevin ÇETİN,

Pamukkale University Faculty of Medicine, Department of Ophthalmology, Denizli, Türkiye

Jan Tjeerd DE FABER

Rotterdam Eye Hospital, Clinic of Pediatric Ophthalmology, Rotterdam, Netherlands

Murat DOĞRU,

Keio University Faculty of Medicine, Department of Ophthalmology, Tokyo, Japan

Ali Hakan DURUKAN,

University of Health Sciences Türkiye, Gülhane Faculty of Medicine Department of Ophthalmology, Ankara, Türkiye

Hayyam KIRATLI,

Hacettepe University Faculty of Medicine, Department of Ophthalmology, Ankara, Türkiye

Tero KIVELÄ,

University of Helsinki, Helsinki University Hospital, Department of Ophthalmology, Helsinki, Finland

Anastasios G.P. KONSTAS,

Aristotle University of Thessaloniki, Department of Ophthalmology, Thessaloniki, Greece

Andrea LEONARDI,

University of Padova, Department of Neuroscience, Unit of Ophthalmology, Padova, Italy

Anat LOEWENSTEIN,

Tel Aviv University Sackler Faculty of Medicine, Department of Ophthalmology, Tel Aviv, Israel

Mehmet Cem MOCAN,

University of Illinois at Chicago, Department of Ophthalmology and Visual Sciences, Illinois, Chicago

Melis PALAMAR ONAY,

Ege University Faculty of Medicine, Department of Ophthalmology, İzmir, Türkiye

Altan Atakan ÖZCAN,

Çukurova University Faculty of Medicine, Department of Ophthalmology, Adana, Türkiye

Özlem ŞAHİN,

Marmara University Faculty of Medicine, Department of Ophthalmology, İstanbul, Türkiye

H. Nida ŞEN,

George Washington University, National Eye Institute, Department of Ophthalmology, Washington, USA

Figen ŞERMET,

Ankara University Faculty of Medicine, Department of Ophthalmology, Ankara, Türkiye

Ebru TOKER,

Marmara University Hospital Faculty of Medicine, Department of Ophthalmology, İstanbul, Türkiye

Şeyda Karadeniz UĞURLU,

İzmir Katip Çelebi University Training and Research Hospital, Department of Ophthalmology, İzmir, Türkiye

Zeliha YAZAR,

University of Health Sciences Türkiye, Ankara Bilkent City Hospital, MHC Building Eye Units Division, Ankara, Türkiye

Nurşen YÜKSEL,

Kocaeli University Faculty of Medicine, Department of Ophthalmology, Kocaeli, Türkiye

The Turkish Journal of Ophthalmology is an official journal of the Turkish Ophthalmological Association.

On Behalf of the Turkish Ophthalmological Association Owner

Hüban ATİLLA

Ankara University Faculty of Medicine, Department of Ophthalmology, Ankara, Türkiye



Publishing House

Molla Gürani Mah. Kaçamak Sokak No: 21,
34093 Fındıkzade-İstanbul-Türkiye

Publisher Certificate Number: 14521

Phone: +90 (530) 177 30 97

E-mail: info@galenos.com.tr

Online Publication Date: June 2025

Publication Type: Local Periodical

International scientific journal published bimonthly.

E-ISSN: 2149-8709

TURKISH JOURNAL OF OPHTHALMOLOGY

TJO



Please refer to the journal's webpage (<https://www.ofthalmoloji.org/>) for "About Us", "Instructions to Authors" and "Ethical Policy".

The editorial and publication process of the Turkish Journal of Ophthalmology are shaped in accordance with the guidelines of ICMJE, WAME, CSE, COPE, EASE, and NISO. The journal adheres to the Principles of Transparency and Best Practice in Scholarly Publishing.

The Turkish Journal of Ophthalmology is indexed in **PubMed/MEDLINE, PubMed Central (PMC), Web of Science-Emerging Sources Citation Index (ESCI), Scopus, TÜBİTAK/ULAKBİM, Directory of Open Access Journals (DOAJ), EBSCO Database, Gale, CINAHL, Proquest, Embase, British Library, Index Copernicus, J-Gate, IdealOnline, Türk Medline, Hinari, GOALI, ARDI, OARE, AGORA, and Turkish Citation Index.**

Issues are published electronically six times a year.

Owner: Hüban ATİLLA on Behalf of the Turkish Ophthalmological Association Owner

Responsible Manager: Banu BOZKURT

CONTENTS

Research Articles

- 112 Comparison of 20% Autologous Platelet-Rich Plasma Versus Conventional Treatment in Moderate to Severe Dry Eye Patients
Shubhi Sachan, Kshama Dwivedi, Satya Prakash Singh, Santosh Kumar, Vinod Kumar Singh; Prayagraj, India
- 120 Comparative Analysis of Automated vs. Expert-Designed Machine Learning Models in Age-Related Macular Degeneration Detection and Classification
Ceren Durmaz Engin, Ufuk Beşenk, Denizcan Özizmirliler, Mustafa Alper Selver; İzmir, Iğdır, Türkiye
- 127 Pattern of RNFL Damage in Early- and Late-Stage Primary Open-Angle Glaucoma Using the Disc Damage Likelihood Scale and Optical Coherence Tomography
Noor Ullah, Rehan Moinuddin Shaikh, Tayyaba Gul Malik; Lahor, Pakistan
- 132 Real-Life Effectiveness and Safety of Selective Laser Trabeculoplasty as Primary, Adjunctive, and Substitutive Therapy
David Oliver-Gutierrez, Osvaldo Guevara-Chavarría, Shirin Djavanmardi, Gloria Segura-Duch, Marta Castany, Stefania Piludu, Carlos A. Arciniegas-Perasso, Elena Ávila, Elena Milla, Susana Duch; Barcelona, Spain
- 141 Gonioscopy-Assisted Transluminal Trabeculotomy versus Bent Ab Interno Needle Goniectomy in Patients with Open-Angle Glaucoma
Ahmet Yücel Üçgöl, Rukiye Kılıç Üçgöl, Zeynep Aktaş; Kırşehir, Ankara, Türkiye

Reviews

- 148 Binocular Approaches in Amblyopia Treatment Based on Dichoptic Stimulation
Demet Yabanoğlu, Hande Taylan Şekeroğlu; Ankara, Türkiye
- 159 Ab Interno Goniotomy/Goniectomy Techniques
Zeynep Aktaş, Syril Dorairaj, Mohamed Sayed, Arsham Sheybani, Ahmet Yücel Üçgöl, Isabella Wagner, Mohamed Khodeiry TX, FL, MO, USA; Abu Dhabi, UAE; Kırşehir, Türkiye

Letters to the Editor

- 171 A Promising Outcome of the Augmented Modified Hummelsheim Procedure in a Challenging Case of Inferior Rectus Hypoplasia
Florentina Priscilia, Anna Puspitasari Bani; Jakarta, Indonesia
- 174 Ptosis Repair by "PEANUTS" MMCR: "Pelın's Easy and Needle Up To Stretch" Müller's Muscle Conjunctival Resection Without the Putterman Clamp
Pelın Kaynak, Gözde Derin Şengün, Can Öztürker, Eda Akgöz; İstanbul, İzmir, Türkiye

AT A GLANCE

2025 Issue 3 at a Glance:

Esteemed colleagues,

In its third issue of 2025, the Turkish Journal of Ophthalmology features five original studies, two reviews, and two letters to the editor.

The prospective interventional study by Sachan et al. titled "Comparison of 20% Autologous Platelet-Rich Plasma Versus Conventional Treatment in Moderate to Severe Dry Eye Patients" included 40 individuals (80 eyes) with moderate to severe dry eye disease. The patients were randomized to the study and control groups (20 each). Those in the study group received 20% autologous platelet-rich plasma eye drops, while the control group received artificial tears as conventional treatment. At the end of three months, there was a significantly greater decrease in ocular surface disease index score in the study group than in the control group, and tear meniscus height, Schirmer test values, and tear breakup time were significantly improved in the study group. The post-treatment improvements in fluorescein staining and impression cytology scores were also significantly greater in the study group compared to the control group. The authors emphasized that autologous platelet-rich plasma therapy is safer and more effective than conventional treatments for moderate to severe symptomatic dry eye disease (See pages 112-119).

In their study titled "Comparative Analysis of Automated vs. Expert-Designed Machine Learning Models in Age-Related Macular Degeneration Detection and Classification", Durmaz Engin et al. aimed to compare the effectiveness of expert-designed machine learning models and code-free automatic machine learning (AutoML) models in detecting age-related macular degeneration (AMD) and distinguishing dry from wet AMD on optical coherence tomography images. The expert-designed model was developed by an AI expert using EfficientNet V2 architecture, while the AutoML model was created by an ophthalmologist using the LobeAI platform, which utilizes transfer learning with ResNet-50 V2. The study compared performance metrics by calculating sensitivity, specificity, accuracy, and F1 scores. The expert-designed model provided 99.67% overall accuracy in classifying all images, and F1 scores were calculated as 0.99 or higher in all binary classifications. On the other hand, while the AutoML model had 89.00% overall accuracy, its F1 scores ranged from 0.86 to 0.90 in binary classifications. The authors stated that although the AutoML model had acceptable performance in recognizing and classifying AMD cases, the expert-designed model had significantly superior performance, emphasizing the importance of the advanced neural network architectures and optimization processes used in expert-developed models (See pages 120-126).

In a cross-sectional study titled "Pattern of RNFL Damage in Early- and Late-Stage Primary Open-Angle Glaucoma Using the Disc Damage Likelihood Scale and Optical Coherence Tomography", UlaIn et al. included 267 eyes of 135 patients aged 18 years and older with suspected or diagnosed glaucoma. After a comprehensive ocular examination, the Disc Damage Likelihood Scale was used for glaucoma staging, with disease severity rated in three zones: green, orange, and red. Retinal nerve fiber layer (RNFL) thickness was measured in the four quadrants using optical coherence tomography, and patterns of RNFL damage were analyzed according to the ISNT rule (inferior>superior>nasal>temporal) and compared between the three groups. The groups differed significantly in terms of mean, inferior, superior, and temporal RNFL thickness, while the difference in nasal RNFL was not significant. While the ISNT rule was found to be the most common pattern among the participants included in the study, progressive loss of this pattern was observed with increased disease severity (See pages 127-131).

In their study titled "Real-Life Effectiveness and Safety of Selective Laser Trabeculoplasty as Primary, Adjunctive, and Substitutive Therapy", Oliver-Gutierrez et al. aimed to evaluate the real-life outcomes of selective laser trabeculoplasty (SLT) in treatment-naïve patients compared to SLT used as adjunctive therapy (AT) and to investigate the potential of SLT to lower intraocular pressure (IOP) and reduce topical medication burden. Patients who had not previously undergone glaucoma surgery or laser treatment received SLT as primary therapy (PT), as AT, or as substitution therapy (ST). Success in the PT and AT groups was defined as $\geq 20\%$ IOP reduction and $\text{IOP} \leq 21$ mmHg in two consecutive follow-ups with the same or fewer drugs without additional glaucoma surgery (including repeat SLT), while success in the ST group was defined as the same or lower IOP with fewer topical medications. The study included a total of 120 eyes of 120 patients with an average follow-up of 32.7 months. The reduction in IOP was greater in the PT group than in the AT group at 24-36 months (22.1% vs. 14.5%, $p=0.039$). Treatment nonresponse was reported in 28.6% of the PT group and 37.0% of the ST group. Success rates were higher in the PT group than in the AT group at 12, 24, and 36 months (47.1% vs. 69.0%, 31.4% vs. 38.8%, and 23.5% vs. 31.1%, respectively). In the ST group, the success rate was 34.2% at 12 months and increased to 38.3% at 24 months (See pages 132-140).

Üçgül et al. conducted a retrospective study titled "Gonioscopy-Assisted Transluminal Trabeculotomy versus Bent Ab Interno Needle Goniectomy in Patients with Open-Angle Glaucoma" comparing the efficacy and safety of gonioscopy-assisted transluminal trabeculotomy (GATT) and bent ab interno needle goniectomy (BANG) in patients with open-angle glaucoma. They evaluated 34 eyes that underwent GATT and 31 eyes that underwent BANG. The preoperative mean IOP was 32.9 ± 6.1 mmHg in the GATT group and 31.8 ± 5.4 mmHg in the BANG group. At the last postoperative follow-up, the mean IOP had decreased to 15.8 ± 4.5 mmHg (51.9% reduction) in the GATT group and 17.9 ± 5.7 mmHg (43.7% reduction) in the BANG group. The rate of complete surgical success was 88.2% for the GATT procedure and 61.3% for the BANG procedure. Early surgical failures were more common

AT A GLANCE

in the BANG group, while the BANG group had fewer early surgical failures and more late surgical failures. While the authors emphasized that GATT surgery provides a larger and more sustainable IOP reduction compared to BANG surgery and has higher surgical success rates, they stated that GATT is a more reliable option in the management of open-angle glaucoma (See pages 141-147).

In their review titled "Binocular Approaches in Amblyopia Treatment Based on Dichoptic Stimulation", Yabanoğlu and Taylan Şekeroğlu systematically examined the literature on dichoptic stimulation techniques that aim to improve binocular function in the treatment of strabismus. Based on various studies in the literature, they explained the basic principles of these treatment methods with emphasis on their results compared to traditional monocular treatment methods. This excellent review provides a comprehensive assessment of the integration of dichoptic treatment approaches into clinical practice and a guiding perspective on the future use of these methods (See pages 148-158).

The second review of the issue was penned by Aktaş et al. and is titled "Ab Interno Goniotomy/Goniectomy Techniques". Minimally invasive glaucoma surgeries (MIGS), which include techniques such as Kahook Dual Blade, BANG, GATT, OMNI, Trabectom, Streamline, and TrabEx+, have made significant advances in the treatment of glaucoma by reducing IOP and improving aqueous humor outflow. These innovative procedures offer effective alternatives to more invasive filtration surgeries by targeting structures such as the trabecular meshwork and Schlemm's canal. MIGS enhances the natural drainage pathways, leading to marked reductions in IOP and reduced dependence on glaucoma medications. Clinical studies show that these MIGS techniques are safe and effective, resulting in fewer complications than traditional surgeries such as trabeculectomy or tube shunt implantation. In addition to providing an overview of the MIGS techniques, this review successfully summarizes the extensive topic of these techniques' evolution, practical applications, and outcomes (See pages 159-170).

In the first letter to the editor in this issue, titled "A Promising Outcome of the Augmented Modified Hummelsheim Procedure in a Challenging Case of Inferior Rectus Hypoplasia", Priscilia and Bani summarized the treatment approach used for a 25-year-old female patient with inferior rectus hypoplasia, a rare pathology (See pages 171-173).

Kaynak et al. share another letter to the editor titled "Ptosis Repair by "PEANUTS" MMCR: "Pelin's Easy and Needle Up To Stretch" Müller's Muscle Conjunctival Resection Without the Putterman Clamp". The authors describe a surgical method that is based on the original Müller muscle-conjunctiva resection technique but uses a 21-gauge needle instead of the Putterman clamp and evaluate the outcomes of three ptosis patients operated using this technique (See pages 174-176).

Covering diagnostic and therapeutic approaches to various ocular pathologies from glaucoma to AMD, from ptosis to dry eye, we hope this issue will provide both scientific and practical guidance for our valued readers.

We would like to thank all the researchers who contributed to this issue. Through your contributions, we hope that each and every issue will feature even richer content.

Respectfully on behalf of the Editorial Board,
Hakan Özdemir, MD